BEST AVAILABLE COPY

Neural precursor and stem cells

Patent number:

EP1529838

Publication date:

2005-05-11

Inventor:

BOSIO ANDREAS DR (DE); CREMER HAROLD (FR);

PENNARTZ SANDRA (DE)

Applicant:

MEMOREC BIOTEC GMBH (DE)

Classification:

- international:

C12N5/06; G01N33/53

- european:

C12N5/06B8P; G01N33/68V2 Application number: EP20030025506 20031106

Priority number(s): EP20030025506 20031106

Cited documents:

EP1354943 XP00227572

XP00222350 XP00103392 XP00902890

Report a data error he

Abstract of EP1529838

A cell population comprising at least 5% neural stem cells, the stem cells being characterized by an expression of ASCT2 or KIAA0152, is new. - Independent claims are also included for the following: - (1) a method for isolating the cell population cited above; - (2) a medicament comprising the above cell population; and - (3) a monoclonal antiboc directed against ASCT2. - ACTIVITY - Neuroprotective; Nootropic; Antiparkinsonian; Cerebroprotective; Vasotropic; No biological data given. - MECHANISM OF ACTION - Cell Therapy.

Data supplied from the esp@cenet database - Worldwide



Europäisches Patentamt

European Patent Office

Office européen des brevets



11) EP 1 529 838 A1

(12)

EUROPÄISCHE PATENTANMELDUNG

- (43) Veröffentlichungstag: 11.05.2005 Patentblatt 2005/19
- (51) Int Cl.7: C12N 5/06, G01N 33/53

- (21) Anmeldenummer: 03025506.1
- (22) Anmeldetag: 06.11.2003
- (84) Benannte Vertragsstaaten:
 AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
 HU IE IT LI LU MC NL PT RO SE SI SK TR
 Benannte Erstreckungsstaaten:
 AL LT LV MK
- (71) Anmelder: Memorec Biotec GmbH 50829 Köln (DE)
- (72) Erfinder:
 - Bosio, Andreas, Dr. 50931 Köln (DE)

- Cremer, Harold
 13400 Marseille (FR)
- Pennartz, Sandra 50670 Köln (DE)
- (74) Vertreter: Schreiber, Christoph, Dr. et al Patentanwälte von Kreisler Selting Werner, Postfach 10 22 41 50462 Köln (DE)
- (54) Neurale Vorläufer- und Stammzellen
- (57) Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Vorläuferzellen sind, die wenigstens einen der in **Liste A** oder **Liste B** aufgeführten Marker aufweisen.

2

Beschreibung

[0001] Die vorliegende Erfindung betrifft Zellpopulationen von neuralen Vorläuferzellen bzw. neuralen Stammzellen sowie Verfahren zur Isolierung entsprechender Zellen.

1

[0002] Der Ausgangspunkt für die Generierung der über tausend verschiedenen neuronalen und glialen Zelltypen des Nervensystems von Vertebraten sind multipotente, neurale Stammzellen des embryonalen Neuroepitheliums (Williams, B. P., Read, J. & Price, J. (1991): The generation of neurons and oligodendrocytes from a common precursor cell. Neuron 7(4), 685-93), (Davis, A. A. & Temple, S. (1994): A self-renewing multipotential stem cell in embryonic rat cerebral cortex. Nature 372(6503), 263-6), (Weiss, S., Dunne, C., Hewson, J., Wohl, C., Wheatley, M., Peterson, A. C. & Reynolds, B. A. (1996): Multipotent CNS stem cells are present in the adult mammalian spinal cord and ventricular neuroaxis. J Neurosci 16(23), 7599-609).

[0003] In den vergangenen Jahren wurde durch verschiedene Arbeitsgruppen gezeigt, dass solche sich selbst erneuernden, multipotenten Vorläuferzellen nicht nur während der Entwicklung, sondern auch im adulten Gehirn zu finden sind (Gage, F. H. (2000): Mammalian neural stem cells. *Science* 287(5457), 1433-8). Vor allem um die lateralen Ventrikel des Vorderhirns findet die Bildung von neuralen Vorläuferzellen lebenslang statt. Diese wandern hauptsächlich, wenn auch nicht exklusiv, in den Bulbus olfaktorius, um dort in GABA-erge Interneurone zu differenzieren.

[0004] Über die genaue Lokalisation der multipotenten Stammzellen, die dieser sekundären Neurogenese zugrunde liegen, wird derzeit noch spekuliert: Johansson et al. beschrieben ependymale Zellen entlang des Lumen der adulten, ventrikulären Zone mit den Eigenschaften multipotenter Stammzellen (Johansson, C. B., Svensson, M., Wallstedt, L., Janson, A. M. & Frisen, J. (1999b): Neural stem cells in the adult human brain. Exp Cell Res 253(2), 733-6), während Doetsch et al. Astrocyten der subventrikulären Zone als multipotente Stammzellen identifizierten (Doetsch, F., Caille, I., Lim, D. A., Garcia-Verdugo, J. M. & Alvarez-Buylla, A. (1999): Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. Cell 97(6), 703-16). Eine absolut eindeutige Identifizierung dieser adulten Stammzellen in vivo ist jedoch bis heute, hauptsächlich mangels geeigneter Marker, nicht gelungen.

[0005] Neben ihrer Bedeutung im olfaktorischen System ist das therapeutische Potential der adulten Stammzellen von besonderem Interesse. Aufgrund ihrer Multipotenz weisen neurale Stammzellen bemerkenswerte Formbarkeit auf und könnten daher durch Zusatz von verschiedenen Faktoren zur Erzeugung verschiedener Neuronentypen eingesetzt werden. Die anschließende Transplantation der so entwickelten spezialisierten Zellen könnte zur Behandlung von neurologischen Krankheiten Alzheimer, Parkinson, Folgen von

Schädel-Hirn-Traumata und Schlaganfall beitragen. Voraussetzung dafür ist die Charakterisierung der verschiedenen, neuralen Differenzierungsstufen sowie die Identifizierung der Faktoren, die die Differenzierungsprogramme der Stammzellen steuern. Gegenüber den embryonalen Stammzellen haben die adulten den Vorteil, dass sie erstens keine abstoßende Immunreaktion auslösen würden, weil sie dem Körper des Patienten entstammen, folglich ihre Transplantation ohne Immunsuppression erfolgen könnte, und zweitens ihre Gewinnung ethisch unbedenklich ist.

[0006] Die Erforschung der Eigenschaften neuraler Stammzellen und embronaler Stammzellen des Menschen ist aus ethischen Aspekten praktisch nicht oder nur sehr eingeschränkt möglich. Daher wurden alle explorativen Arbeiten ausgehend von Mäusen und Mauszellen durchgeführt. Wie bereits beschrieben war die Isolierung von neuralen Stammzellen bisher nicht möglich, da dieser Zelltyp nicht eindeutig charakterisiert war und keine geeigneten Marker zur Identifizierung und Anreicherung zur Verfügung standen.

[0007] Aufgabe der vorliegenden Erfindung war es daher Verfahren zu entwickeln, die eine Isolation von neuralen Vorläuferzellen und neuralen Stammzellen erlauben und entsprechende Zeilpopulation, enthaltend diese Zelle bereitzustellen.

[0008] Erfindungsgemäß wird die Aufgabe gelöst durch die Identifizierung von Markern, die entsprechende Zellen aufweisen.

30 [0009] Marker ist ein Gen, das mit Hilfe der Serial Analysis of Genexpression (SAGE) in entsprechenden Zellen gefunden wird.

[0010] Methodisch beruht SAGE auf der Isolierung von 14 bp großen DNA Fragmenten (Tags), die jeweils charakteristisch für eine mRNA-Spezies sind. Die Tags. repräsentativ für alle in der zu untersuchenden Zelle vorliegenden mRNA Moleküle, werden zu langen Polymeren verbunden, die im letzten Schritt der Methode sequenziert werden. Die Frequenz, mit der ein Tag sequenziert wird, ist direkt proportional zur Kopienzahl der mRNA-Moleküle im untersuchten Ausgangsmaterial (Velculescu, V. E., Zhang, L., Vogelstein, B. & Kinzler, K. W. (1995): Serial analysis of gene expression. Science 270(5235), 484-7). Durch die computerunterstützte Auswertung der Sequenzdaten entsteht ein digitales Expressionsprofil, das beliebig oft und ohne zusätzliche Laborarbeit mit Expressionsprofilen anderer Gewebe verglichen werden kann (Meta-Analyse).

[0011] Den so identifizierten Gene sind eindeutigen Nummern zugeordnet, die beispielsweise als SAGEmap von National Center for Biotechnology Information (NCBI) bereitgestellt werden (www.ncbi.nlm.nin.gov/SAGE).

[0012] Gegenstand der Erfindung sind zum einen Zellpopulationen, bei denen mindestens 5% der Zellen neurale Vorläuferzellen sind, die wenigstens einen der in Liste A oder Liste B aufgeführten Marker aufweisen. [0013] Bevorzugt weisen entsprechende neurale Vor-

läuferzellen wenigstens zwei, drei, vier oder fünf der in Liste A oder B aufgeführten Marker auf.

3

[0014] In bevorzugten Ausführungsformen weisen entsprechende neurale Vorläuferzellen keinen der in Liste C aufgeführten Marker auf.

[0015] Bevorzugt ist der Gehalt an neuralen Vorläuferzellen in der Zellpopulation hoch, d.h. mindestens 10%, bevorzugt mindestens 25%, noch mehr bevorzugt mehr als 50% und am meisten bevorzugt über 90%.

[0016] Entsprechende neurale Vorläuferzellen sind vorzugsweise aus Hirngewebe erhältlich.

[0017] In einer Ausführungsform handelt es sich dabei um eine murine Zellpopulation.

[0018] Gegenstand der Erfindung ist auch ein Verfahren zur Isolierung einer entsprechenden Zellpopulation mit folgenden Schritten:

entweder

- Entnahme einer Probe aus dem Hirn
- Isolieren der neuralen Vorläuferzellen unter Ver- 20 wendung der angegebenen Marker

oder

- Differenzierung von embryonalen Stammzellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- Differenzierung von adulten, neuralen Stammzellen zu neuralen Vorläuferzellen.
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- Differenzierung von immortalisierten Zellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker.

[0019] "Unter Verwendung der angegebenen Marker" bedeutet, dass die Zellen isoliert werden, die positiv für mindestens einen der Marker aus der Liste A und B sind, wobei mehrere positive Marker und die Abwesenheit von Markern der Liste C bevorzugt werden. Die Isolierung kann beispielsweise durch FACS Analyse erfolierung kann beispielsweise durch FACS Analyse er

gen. Die durch die Verfahren erhältlichen Zellen sind ebenfalls Gegenstand der Erfindung.

[0020] Ein weiterer Gegenstand der Erfindung ist die Verwendung mindestens eines Markers ausgewählt

aus der Liste A oder Liste B zu identifizierung oder isolierung von neuralen Vorläuferzellen.

[0021] Gegenstand ist weiterhin ein Antikörper gegen einen Marker aus der Liste A, B oder C, ein Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste A, B oder C sowie ein Arzneimittel enthaltend die erfindungsgemäße Zellpopulation.

[0022] Solche Arzneimittel könnten wie oben dargestellt zur Behandlung von neurologischen Krankheiten wie Alzheimer, Parkinson, Folgen von Schädelhirntraumata oder Schlaganfall eingesetzt werden.

[0023] Ein weiterer Gegenstand ist eine Zellpopulation, bei der mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens einen der in Liste D oder Liste E aufgeführten Marker aufweisen.

[0024] Vorzugsweise weisen entsprechende neurale Stammzellen mindestens zwei, bevorzugt mindestens drei, mindestens vier und noch mehr bevorzugt mindestens fünf der in Liste D oder Liste E aufgeführten Marker auf.

[0025] In besonders bevorzugten Ausführungsformen weisen entsprechende neurale Stammzellen keinen der in Liste A oder Liste C aufgeführten Marker auf. [0026] Der Gehalt an neuralen Stammzellen in der Zellpopulation ist möglichst hoch, bevorzugt mindestes 10%, mehr bevorzugt mindestes 25%, mindestens 50%, und am meisten bevorzugt mindestens 90%.

[0027] Entsprechende Zellpopulation sind aus Hirngewebe erhältlich. In einer Ausführungsform handelt es sich um eine murine Zellpopulation.

[0028] Gegenstand ist weiterhin ein Verfahren zur Isolierung der Zellpopulation. Dies ist erhältlich entweder durch

- Entnahme einer Probe aus dem Hirn
 - Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

45 oder

- Differenzierung von embryonalen Stammzellen zu neuralen Stammzellen,
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Stammzellen.
 - Isolieren der neuralen Stammzellen unter Verwen-

dung der angegebenen Marker

oder

- De-Differenzierung von adulten, neuralen Vorfäuferzellen zu neuralen Stammzellen,
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- Differenzierung von immortalisierten Zellen zu neuralen Stammzellen,
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker.

[0029] Die Isolierung erfolgt wie oben bei den neuralen Vorläuferzellen angegeben. Auch die auf diesem Wege erhältlichen neuralen Stammzellen sind Gegenstand der Erfindung.

[0030] Gegenstand der Erfindung ist weiterhin ein Antikörper gegen einen Marker aus der Liste D, E, ein Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste D, E, A oder C sowie ein Arzneimittel enthaltend die erfindungsgemäße Zellpopulation.

[0031] Solche Arzneimitteln können wie dargestellt zur Behandlung von neuronalen Krankheiten wie Alzheimer, Parkinson, Folgen von Schädelhirntraumata oder Schlaganfall eingesetzt werden.

Beispiele

A. Isolierung von embryonaler Stammzellen

[0032] Murine embryonale Stammzellen proliferieren klonal in vitro und sind aus diesem Grunde in großer Menge und hochreiner Form isolierbar. Nach dem Stand der Technik werden diese in Anwesenheit von LIF auf primären embryonalen Fibroblasten gehalten und regelmäßig durch die Generierung von hochgradig keimbahnkompetenten chimären Mäusen auf ihre Qualität überprüft. Unter normalen Kulturbedingungen beträgt das Verhältnis ES-Zellen zu kontaminierenden Fibroblasten etwa 200:1. Um auch diese minoritäre Komponente zu eliminieren, wurden die ES-Zellen vor der RNA-Päparation für zwei Passagen (vier Tage) auf gelatinisierten Kulturplatten bei erhöhter LIF-Konzentration gehalten. Dies ermöglicht eine Reduktion der kontaminierenden Fibroblasten auf etwa 0,01% der Gesamtpopulation.

B. Isolierung von neuronalen Vorläuferzellen aus dem adulten Mausgehirn.

[0033] In der subventrikulären Zone des adulten Vor-

derhirns von Vertebraten werden permanent große Mengen von neuralen Vorläuferzellen gebildet (wahrscheinlich < 50000 Zellen/ Tag). Diese Zellen benutzen einen präzise definierten Migrationsweg und eine spezielle Form der Translokation (*Chain migration*) um in den Bulbus olfaktorius zu gelangen. Im Bulbus olfaktorius angelangt differenzieren diese Vorläuferzellen normalerweise in inhibitorische (GABA-erge) Interneurone. Unter bestimmten experimentellen Bedingungen wurde ihre Differenzierung in Oligodendrozyten und Astrozy-

6

ten gezeigt.

[0034] Neurale Vorläufer, die einen Differenzierungszustand zwischen einer neuralen Stammzelle und einem terminal differenzierten Neuron repräsentieren, exprimieren spezifisch eine Form des neuralen Zelladhäsionsmoleküls NCAM, die eine spezielle post-translationelle Modifikation aufweist. Diese Modifikation besteht aus der Glykosylierung des Proteins mit a-2,8 verknüpfter Polysialylsäure (PSA). Ein spezifischer Antikörper gegen dieses Glykoepitop (Chazal et al., 2000) erlaubte die hochreine Isolierung der Zielpopulation aus dissozierten Vorderhirngewebe durch FACS (Fluorescence Activated Cell Sorting).

C. Molekulargenetische Analyse

[0035] Embryonale Stammzellen und neuronale Vorläuferzellen wurden in einem genomweiten Screen mit der Methode SAGE (Serial Analysis of Gene Expression) analysiert.

[0036] Die Genexpressionsprofile der beiden Zell-Populationen wurden unter Anwendung bioinformatischer Verfahrensweisen mit Maus-Hirn-SAGE-Datenbanken verglichen, um molekulare Marker zu identifizieren, die charakteristisch für embryonale Stammzellen und neuronale Vorläuferzellen sind.

[0037] Mit Hilfe der Microarray technologie wurde die Expression der Gene bestätigt.

[0038] Durch in situ-Hybridisierung in Maushirn und an embryonalen Stammzellen wurde die zelluläre Lokalisation einiger der identifizierten Gene bestimmt. Diese Ergebnisse belegen, dass spezifische Markergene identifiziert werden konnten.

Liste A: Positivmarker neurale Vorläuferzellen (1.) und Negativmarker 2 neurale Stammzellen;

ES-Zellen -; PSA-NCAM +; Adult brain -

50 [0039]

	Mm.8884	nuclear factor of kappa light chain gene
		enhancer in B-cells inhibitor, alpha
	[°] Mm.8180	lymphocyte antigen 6 complex, locus A
5	Mm.6238	SRY-box containing gene 11
	Mm.517	(Manual) Manic fringe protein, putative
		secreted glycosyltransferase, notch
		modulator

7

Mm.4919 Mm.4727	DNA segment, human D4S114 seizure related gene 6			tivmarker neurale Vorläuferzeilen (2.);
Mm.45769 Mm.44490	ESTs RIKEN cDNA 6330415M09 gene		ES-Zellen -/+	; PSA-NCAM +; Adult brain -
Mm.42948	peroxiredoxin 2	5	[0040]	
Mm.4022 Mm.3940	RIKEN cDNA 1110033C18 gene lethal giant larvae homolog		Mm.911	high mobility group nucleosomal bin-
Mm.37835	ribosomal protein L7			ding domain 2
Mm.3779	RIKEN cDNA 2300006C11 gene		Mm.89136	H3 histone, family 3A
Mm.340	high mobility group box 3	10		fatty acid binding protein 5, epidermal
Mm.32902	ESTs, Weakly similar to S26689 hypothetical protein had a mayor		Mm.7286	C-terminal binding protein 1
Mm.3268	thetical protein hc1 - mouse ubiquitin-conjugating enzyme E2I		Mm.7141 Mm.6840	proliferating cell nuclear antigen RIKEN cDNA 5730507C05 gene
Mm.31436	myeloid ecotropic viral integration site-		Mm.6787	splicing factor, arginine/serine-rich 3
	related gene 1	15		(SRp20)
Mm.297	actin, beta, cytoplasmic		Mm.6417	CD24a antigen
Mm.29558	expressed sequence Al426163		Mm.6343	nucleophosmin 1
Mm.29014	T-cell lymphoma invasion and metasta-		Mm.482	Jun oncogene
	sis 2		Mm.43871	expressed sequence AW046487
Mm.28842	chloride channel 3	20	Mm.43213	RIKEN cDNA 9030402K04 gene
Mm.28824	Mus musculus, clone IMAGE:4504748,		Mm.42767	ribosomal protein S17
Mm.28275	mRNA RNA binding motif protein, X chromoso-		Mm.4269 Mm.40715	transcription factor 4 RIKEN cDNA 1110038H03 gene
WIIII.20273	me		Mm.40715	RIKEN cDNA 1110038H03 gene
Mm.28149	RIKEN cDNA 3110003A17 gene	25	Mm.4071	laminin receptor 1 (67kD, ribosomal pro-
Mm.28148	chromobox homolog 3 (Drosophila HP1			tein SA)
	gamma)		Mm.4025	nuclear factor I/B
Mm.27816	hexosaminidase B		Mm.372	ribosomal protein S26
Mm.2769	MARCKS-like protein		Mm.3487	ribosomal protein L30
Mm.22171	calponin 3, acidic	30	Mm.3381	ribosomal protein S8
Mm.220923 Mm.21740	RIKEN cDNA 6530406007 gene heterogeneous nuclear ribonucleopro-		Mm.31051	RIKEN cDNA 2610003J05 gene
WIIII.21740	tein H1		Mm.30120 Mm.30011	ribosomal protein S27-like ribosomal protein S23
Mm.206085	expressed sequence Al854782		Mm.29911	RIKEN cDNA 3200001M24 gene
Mm.205996	EST AA087124	35	Mm.2966	isocitrate dehydrogenase 2 (NADP+),
Mm.200858	RIKEN cDNA 2410129E14 gene			mitochondrial
Mm.199500	expressed sequence Al844617		Mm.29580	superiorcervical ganglia, neural specific
Mm.195901	ribosomal protein L35a			10
Mm.194965	EST		Mm.2958	expressed sequence Al843786
Mm.19101	DEAD (aspartate-glutamate-alanine-aspartate) box polypeptide 5	40	Mm.28985 Mm.28869	ribosomal protein L27 ESTs
Mm.19016	drebrin 1		Mm.27927	heterogeneous nuclear ribonucleopro-
Mm.18789	SRY-box containing gene 4		141111.27327	tein A1
Mm.186740	ESTs		Mm.27669	small nuclear ribonucleoprotein E
Mm.18516	H3 histone, family 3B	45	Mm.2756.	high mobility group nucleosomal bin-
Mm.181959	early growth response 1			ding domain 1
Mm.181847	prefoldin 5		Mm.27141	Rac GTPase-activating protein 1
Mm.16421	high mobility group box 1		Mm.2591	RNA binding motif protein 3
Mm. 15534	interleukin 1 alpha	50	Mm.24083	Mus musculus, Similar to TAR DNA bin-
Mm.13725 Mm.12871	Paneth cell enhanced expression doublecortin	50		ding protein, clone MGC: 19284 IMAGE:4016437, mRNA, complete cds
Mm.127662	ESTs		Mm.219668	RIKEN cDNA 2610209F03 gene
Mm.12412	Mus musculus, Similar to RIKEN cDNA 2810407E23 gene, clone IMAGE:		Mm.21841	splicing factor, arginine/serine-rich 2 (SC-35)
	4489006, mRNA, partial cds	55	Mm.218240	Mus musculus, clone IMAGE:5342828, mRNA, partial cds
			Mm.21740	heterogeneous nuclear ribonucleoprotein H1

EP	1	529	838	A 1

	(Manual) 60S ribosomal protein L32		Mm.6660	small inducible cytokine A27
Mm.2115	(RPL32) heterogeneous nuclear ribonucleopro-		Mm.6586	Mus musculus, clone MGC:6299 IMAGE: 2654341, mRNA, complete cds
	tein U		Mm.6565	FK506 binding protein 8 (38 kDa)
Mm.196611	synapsin I	5	Mm.65337	Mus musculus, clone MGC:28924
Mm.19187	prothymosin alpha			IMAGE:3481738, mRNA, complete cds
Mm.18789	SRY-box containing gene 4		Mm.648	prion protein
Mm.186499	ESTs, Weakly similar to immunoglobulin		Mm.638	ESTs
	superfamily containing leucinerich re-		Mm.544	phosphoprotein enriched in astrocytes 15
	peat	10	Mm.5264	ESTs, Highly similar to FEZ1_RAT FA-
Mm.18516	H3 histone, family 3B			SCICULATION AND ELONGATION
Mm.180873	RIKEN cDNA 2510019J09 gene			PROTEIN ZETA 1 (ZYGIN I)
Mm.1775	hematological and neurological expres-		Mm.5259	(Manual assignment) probably myelin-
	sed sequence 1			associated oligodendrocyte basic protein
Mm.1703	tubulin, beta 5	15		MOBP
Mm.16775	ribosomal protein S24		Mm.5249	copine 6
Mm.16767	heterogeneous nuclear ribonucleopro-		Mm.52	RIKEN cDNA 1810033A19 gene
	tein A2/B1		Mm.5195	complexin 1
Mm.16596	B-cell translocation gene 1, anti-prolife-		Mm.5153	neurotensin receptor 2
	rative	20	Mm.5023	Purkinje cell protein 4
Mm.148973	RIKEN cDNA 3010025E17 gene		Mm.4923	ESTs
Mm.142872	heterogeneous nuclear ribonucleopro-		Mm.4921	glutamate receptor, ionotropic, AMPA2
	tein K			(alpha 2)
Mm.142729	thymosin, beta 4, X chromosome		Mm.4920	glutamate receptor, ionotropic, AMPA1
Mm.140380	ribosomal protein L23	25		(alpha 1)
Mm.140	protein phosphatase 1, regulatory (inhi-		Mm.4870	synaptosomal-associated protein, 91
	bitor) subunit 14B			kDa
Mm. 12858	eukaryotic translation initiation factor		Mm.4857	calcium/calmodulin-dependent protein
	4A1			kinase II, beta
		30	Mm.4762	kinesin heavy chain member 1A
-	ativmarker 1 neurale Stammzellen und		Mm.4705	(Manual) probably in far 3'-UTR of com-
Negativmar	ker neurale Vorläuferzellen;			plexin-2 cDNA
F0 7 "	Des 116.11		Mm.46764	RIKEN cDNA 4833409J18 gene
ES-Zellen -;	PSA-NCAM -; Adult brain +		Mm.4657	amyloid beta (A4) precursor protein-bin-
[0044]		35	NA:- 4054	ding, family A, member 2
[0041]			Mm.4651	
M== 00				kinesin-associated protein 3
	protogomo (procomo mocronoin) eubu		Mm.45951	RIKEN cDNA 1200016B17 gene
Mm.98	proteasome (prosome, macropain) subu-			RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 po-
	nit, beta type 6	40	Mm.45951 Mm.4550	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 po- lypeptide
Mm.9745	nit, beta type 6 lactate dehydrogenase 2, B chain	40	Mm.45951	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 po- lypeptide ATPase, Na+/K+ transporting, beta 1 po-
	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui-	40	Mm.45951 Mm.4550 Mm.4550	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 po- lypeptide ATPase, Na+/K+ transporting, beta 1 po- lypeptide
Mm.9745 Mm.970	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous	40	Mm.45951 Mm.4550	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 po- lypeptide ATPase, Na+/K+ transporting, beta 1 po- lypeptide NADH dehydrogenase (ubiquinone) 1
Mm.9745 Mm.970 Mm.891	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2	40	Mm.45951 Mm.4550 Mm.4550 Mm.4537	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9
Mm.9745 Mm.970	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-CI cotrans-	40 45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene
Mm.9745 Mm.970 Mm.891 Mm.88833	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds		Mm.45951 Mm.4550 Mm.4550 Mm.4537	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen		Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa
Mm.9745 Mm.970 Mm.891 Mm.88833	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene		Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen		Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6		Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ trans-
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261	45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen	45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC:
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat se- quence-containing transcript aldolase 3, C isoform	45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, comple-
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat se- quence-containing transcript	45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107 Mm.44101	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-Cl cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat se- quence-containing transcript aldolase 3, C isoform	45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107 Mm.44101	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123 Mm.7729 Mm.7420 Mm.7363 Mm.726	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-CI cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat se- quence-containing transcript aldolase 3, C isoform tubulin, beta 4 beta-spectrin 3 basigin	45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107 Mm.44101 Mm.4383 Mm.43786	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1 cytochrome c oxidase, subunit VIIc
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.806 Mm.80123 Mm.7729 Mm.7420 Mm.7363 Mm.726 Mm.7089	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-CI cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat se- quence-containing transcript aldolase 3, C isoform tubulin, beta 4 beta-spectrin 3 basigin necdin	45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107 Mm.44101 Mm.4383 Mm.43786 Mm.43749 Mm.43721 Mm.43587	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1 cytochrome c oxidase, subunit VIIc RIKEN cDNA 3100001N19 gene small nuclear ribonucleoprotein N hippocalcin
Mm.9745 Mm.970 Mm.891 Mm.88833 Mm.87027 Mm.8688 Mm.86654 Mm.848 Mm.806 Mm.80123 Mm.7729 Mm.7420 Mm.7363 Mm.726	nit, beta type 6 lactate dehydrogenase 2, B chain creatine kinase, mitochondrial 1, ubiqui- tous kinesin family member C2 Mus musculus strain ILS K-CI cotrans- porter (Slc12a5) mRNA, complete cds BM88 antigen RIKEN cDNA 0610011B04 gene microtubule-associated protein 6 testis expressed gene 261 CD 81 antigen ESTs, Weakly similar to simple repeat se- quence-containing transcript aldolase 3, C isoform tubulin, beta 4 beta-spectrin 3 basigin	45	Mm.45951 Mm.4550 Mm.4550 Mm.4537 Mm.44355 Mm.4435 Mm.44244 Mm.44107 Mm.44101 Mm.4383 Mm.43786 Mm.43749 Mm.43721	RIKEN cDNA 1200016B17 gene ATPase, Na+/K+ transporting, beta 1 polypeptide ATPase, Na+/K+ transporting, beta 1 polypeptide NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 9 RIKEN cDNA 6430514L14 gene synaptosomal-associated protein, 25 kDa open reading frame 12 ESTs Mus musculus, ATPase, Na+K+ transporting, alpha 3 subunit, clone MGC: 27631 IMAGE:4506376, mRNA, complete cds myc box dependent interacting protein 1 cytochrome c oxidase, subunit VIIc RIKEN cDNA 3100001N19 gene small nuclear ribonucleoprotein N

	EΡ	1	529	838	A1
--	----	---	-----	-----	----

	lypeptide 1		Mm.3974	ubiquitin specific protease 4 (proto-onco-
Mm.4339	laminin, alpha 5			gene)
Mm.43330	RIKEN cDNA 0610025G13 gene		Mm.39548	expressed sequence AI839779
Mm.43278	olfactomedin 1		Mm.3951	thymus cell antigen 1, theta
Mm.43278	olfactomedin 1	5	Mm.3915	myelin-associated oligodendrocytic ba-
Mm.4296	synovial sarcoma translocation, Chromo-			sic protein
	some 18		Mm.39040	myelin and lymphocyte protein, T-cell dif-
Mm.42949	RIKEN cDNA 1110012005 gene			ferentiation protein
Mm.42948	peroxiredoxin 2		Mm.38994	RIKEN cDNA 2600001N01 gene
Mm.42829	selenoprotein W, muscle 1	10	Mm.38993	calsyntenin 1
Mm.4266	integral membrane protein 2B		Mm.38551	calcium binding protein 1
Mm.4266	integral membrane protein 2B		Mm.38469	amyloid beta (A4) precursor protein-bin-
Mm.4263	cystatin C			ding, family B, member 1
Mm.425	histidine triad nucleotide binding protein		Mm.38438	RIKEN cDNA 1200009K17 gene
Mm.42255	ATPase, Ca++ transporting, cardiac	15	Mm.38421	(Manual assignment) ATPase, Na+K+
	muscle, slow twitch 2			transporting, alpha polypeptide
Mm.41926	NADH dehydrogenase (ubiquinone) 1 al-		Mm.38421	(Manual assignment) ATPase, Na+K+
	pha subcomplex, 4			transporting, alpha polypeptide
Mm.41925	RIKEN cDNA 1810034B16 gene		Mm.3840	flotillin 2
Mm.41918	RIKEN cDNA 1110063G11 gene	20	Mm.38248	sialyltransferase 9 (CMP-NeuAc:lacto-
Mm.41911	cytochrome P450, 46 (cholesterol 24-hy-			sylceramide alpha-2,3-sialyltransferase)
	droxylase)		Mm.38036	ESTs, Moderately similar to
Mm.41893	RIKEN cDNA 6330408G06 gene			NX1A_MOUSE_2
Mm.41791	glycoprotein m6b		Mm.38036	ESTs, Moderately similar to
Mm.41752	expressed sequence Al847934	25		NX1A MOUSE 2
Mm.41735	RIKEN cDNA 2300004C15 gene		Mm.37462	ESTs, Weakly similar to CA11 RAT COL-
Mm.41719	RIKEN cDNA 2610507A21 gene		14111.57402	LAGEN ALPHA 1(I) CHAIN
Mm.41711	Mus musculus, clone IMAGE:3499845,		Mm.37214	transferrin
191111.477717	mRNA, partial cds		Mm.36275	DNA segment, Chr 11, Brigham & Wo-
Mm.41694	ESTs	30	WIIII.30275	men's Genetics 0517 expressed
Mm.41692	ESTs, Weakly similar to F59F4.2.p	50	Mm.3624	•
Mm.41642	regulator of G-protein signaling 4		Mm.35837	guanylate kinase 1
Mm.41630	RIKEN cDNA 0710001E10 gene			RIKEN cDNA 2510006D16 gene
Mm.41604	ESTs, Weakly similar to VAV3_MOUSE		Mm.35837	RIKEN cDNA 2510006D16 gene
WIII.41004	VAV-3 PROTEIN	35	Mm.3544	calcium channel, voltage-dependent, be- ta 3 subunit
Mm.41603	expressed sequence Al891706		Mm.35439	secreted acidic cysteine rich glycoprotein
Mm.41603	expressed sequence Al891706		Mm.35270	Ly6/neurotoxin 1
Mm.41602	RIKEN cDNA 3110050007 gene		Mm.3479	ATPase, H+ transporting, lysosomal
Mm.41602	RIKEN cDNA 3110050007 gene			21kDa, V0 subunit B
Mm.4137	chromogranin A	40	Mm.34695	actin related protein 2/3 complex, subunit
Mm.41354	ESTs		1111110-1000	1A (41 kDa)
Mm.41277	RIKEN cDNA 1110020M21 gene		Mm.34246	calmodulin 1
Mm.41248	ESTs		Mm.3363	prosaposin
Mm.41190	RIKEN cDNA 1700112L09 gene		Mm.3360	tyrosine 3-monooxygenase/tryptophan
Mm.40863	expressed sequence AW049870	45	141111.0000	5-monooxygenase activation protein, ze-
Mm.40738	RIKEN cDNA 2900072M03 gene			ta polypeptide
Mm.40621	ESTs, Moderately similar to		Mm.33117	ESTs
	Y552_HUMAN HYPOTHETICAL PRO-		Mm.3308	tyrosine 3-monooxygenase/tryptophan
	TEIN KIAA0552		141111.0000	5-monooxygenase activation protein, eta
Mm.40472	expressed sequence Al835002	50		polypeptide
Mm.40443	RIKEN cDNA 4930488B01 gene		Mm.3292	glutamate receptor, ionotropic, NMDA1
Mm.40124	phosphodiesterase 10A		14011.0232	(zeta 1)
Mm.40059	ESTs, Weakly similar to SP62 MOUSE		Mm.3229	ribosomal protein L26
	SPLICEOSOME ASSOCIATED PROTE-		Mm.32191	gamma-aminobutyric acid (GABA-B) re-
	IN 62	55	141111.02 13 1	ceptor, 1
Mm.39857	RIKEN cDNA 2900074L19 gene		Mm.31395	carboxypeptidase E
Mm.39803	expressed sequence Al841080		Mm.3123	cornichon-like (Drosophila)
Mm.39752	RIKEN cDNA 2900041A09 gene		Mm.31025	RIKEN cDNA 2310015K15 gene
	obian 200004 inos gene		141111.51025	THICK ODIAN 20100 ISK IS gelle

Mm.29344

Mm.29330

tumor differentially expressed 1, like

expressed sequence Al853543

EP ·	1	529	838	A.
------	---	-----	-----	----

13 14 Mm.30412 RIKEN cDNA 5430400P17 gene Mm.29230 RIKEN cDNA 1500017E18 gene Mm.30355 (Manual) KIF5A Neuronal Kinesin heavy Mm.29227 RIKEN cDNA 2300002D11 gene Mm.29205 bruno-like 4, RNA binding protein (Droso-Mm.30266 hemoglobin, beta adult major chain phila) Mm.30266 hemoglobin, beta adult major chain Mm.29205 bruno-like 4, RNA binding protein (Droso-Mm.30206 ATPase, H+ transporting, lysosomal phila) 34kD, V1 subunit D Mm.2918 megakaryocyte-associated tyrosine kiprotease, serine, 11 (Igf binding) Mm.30156 Mm.30155 ATPase, H+ transporting, lysosomal Mm.29141 RIKEN cDNA 0710008N11 gene 16kD, V0 subunit C 10 Mm.29124 phosphatidic acid phosphatase type 2B Mm.30150 RIKEN cDNA 1010001M12 gene Mm.29075 (Manual) Reticulon 1 protein, major inter-Mm.30126 membrane interacting protein of RGS16 Mm.30085 aldo-keto reductase family 1, member A4 Mm.29027 SPARC-like 1 (mast9, hevin) (aldehyde reductase) Mm.29027 SPARC-like 1 (mast9, hevin) Mm.30072 cytochrome c oxidase subunit VIIa poly-Mm.2902 protein tyrosine phosphatase, receptorpeptide 2-like Mm.30059 myristoylated alanine rich protein kinase Mm.28955 RIKEN cDNA 4930570C03 gene C substrate Mm.28650 RAB6, member RAS oncogene family Mm.29976 septin 5 Mm.28650 RAB6, member RAS oncogene family Mm.29965 RIKEN cDNA 2410104119 gene Mm.28643 vesicle-associated membrane protein 2 Mm.29947 serine/threonine kinase 11 Mm.28561 protein kinase C, zeta Mm.29939 RIKEN cDNA 1010001N11 gene Mm.28518 type I transmembrane protein Fn14 Mm.29937 (Manual assignment) polymorphism of Mm.28357 microtubule-associated protein 1 light Mm.29937 ESTs, Weakly similar to predicted using Genefinder 25 Mm.2815 RIKEN cDNA 1110021H02 gene Mm.29921 RAS protein-specific guanine nucleotide-Mm.28107 ectonucleotide pyrophosphatase/phosreleasing factor 1 phodiesterase 2 Mm.2992 (Manual assignment) MBP myelin basic Mm.28058 NADH dehydrogenase (ubiquinone) 1 protein beta subcomplex 5 Mm.29870 integral membrane protein 3 Mm.27886 RIKEN cDNA 2410011G03 gene Mm.29867 NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 Mm.27608 Mus musculus, Similar to chromosome Mm.29857 (Manual) Neurogranin 9 open reading frame 16, clone MGC: Mm.29852 Mus musculus, clone IMAGE:5102170, 19388 IMAGE:2812475, mRNA, commRNA, partial cds 35 plete cds Mm.29846 Mus musculus, Similar to NDRG family, Mm.2755 calbindin 2 member 4, clone MGC:7067 IMAGE: Mm.27499 RIKEN cDNA 2010004E11 gene 3156802, mRNA, complete cds Mm.27407 RecQ protein-like Mm.29842 NADH dehydrogenase flavoprotein 1 Mm.27256 discs, large homolog 4 (Drosophila) Mm.29823 microsomal glutathione S-transferase 3 40 Mm.2720 mitogen activated protein kinase 8 in-Mm.29807 ubiquitin carboxy-terminal hydrolase L1 teracting protein Mm.29807 ubiquitin carboxy-terminal hydrolase L1 Mm.27114 RIKEN cDNA 0610043B10 gene Mm.29771 ATPase, H+ transporting, lysosomal Mm.27087 RIKEN cDNA 2010012C24 gene 70kD, V1 subunit A, isoform 1 Mm.27005 visinin-like 1 Mm.29717 45 3-monooxgenase/tryptophan 5-monoox-Mm.26633 PH domain containing protein in retina 1 genase activation protein, gamma poly-Mm.26633 PH domain containing protein in retina 1 peptide phosphofructokinase, muscle Mm.26550 Mm.29711 adrenergic receptor kinase, beta 1 Mm.2645 eukaryotic translation elongation factor Mm.297 actin, beta, cytoplasmic 1 alpha 2 Mm.29633 RIKEN cDNA 1810008021 gene 50 Mm.2635 pyruvate kinase 3 Mm.29600 Mus musculus, clone IMAGE:3964267, Mm.2619 cholecystokinin mRNA Mm.25849 RIKEN cDNA 2010003014 gene Mm.2948 H2-K region expressed gene 2 Mm.25738 RIKEN cDNA 2900002P20 gene Mm.29477 SCAN domain-containing 1 Mm.25228 ring finger protein 11 Mm.29415 RIKEN cDNA 1810011001 gene Mm.25203 NCK-associated protein 1 Mm.29362 expressed sequence Al414999 Mm.2496 internexin neuronal intermediate fila-

Mm.24482

ment protein, alpha

RIKEN cDNA 5730460C18 gene

15

Mm.2446	synaptotagmin 4			pha o
Mm.24376	Mus musculus mRNA for calsyntenin-3		Mm.20964	guanine nucleotide binding protein, al-
	(Cs3 gene)		141111.20304	pha o
Mm,2411	Ras-GTPase-activating protein (GAP		Mm.2082	apolipoprotein D
	<120>) SH3-domain binding protein 2	5	Mm.206218	Mus musculus, Similar to hypothetical
Mm.24092	N-ethylmaleimide sensitive fusion pro-	_	141111.200210	protein FLJ22237, clone MGC:27683
***************************************	tein			IMAGE:4913322, mRNA, complete cds
Mm.24092	N-ethylmaleimide sensitive fusion pro-		Mm.2060	RIKEN cDNA 2900010105 gene
	tein		Mm.20472	vertebrate homolog of C. elegans Lin-7
Mm.2400	glutathione peroxidase 4	10		type 2
Mm.2397	synaptophysin		Mm.203939	expressed sequence Al256814
Mm.23826	phosphotyrosyl phosphatase activator		Mm.203924	expressed sequence AW259572
Mm.2381	amyloid beta (A4) precursor-like protein		Mm.203921	expressed sequence Al850305
	1		Mm.202728	expressed sequence Al447901
Mm.2338	glutamine synthetase	15	Mm.202696	expressed sequence AA409221
Mm.2338	glutamine synthetase		Mm.201729	expressed sequence Al426007
Mm.2326	macrophage migration inhibitory factor		Mm.2011	glutathione S-transferase, mu 1
Mm.2319	Scgn10 like-protein		Mm.200858	RIKEN cDNA 2410129E14 gene
Mm.23023	RIKEN cDNA 1500009C09 gene		Mm.200843	synuclein, beta
Mm.23002	RIKEN cDNA 5330410G16 gene	20	Mm.200817	expressed sequence AW124717
Mm.22699	selenoprotein P, plasma, 1		Mm.200817	expressed sequence AW124717
Mm.22637	RIKEN cDNA 0910001L24 gene		Mm.200806	(Manual) no clear assignment, probably
Mm.22597	RIKEN cDNA 2310042E05 gene			non-coding (but spliced) RNA gene
Mm.22473	Rab acceptor 1 (prenylated)		Mm.200511	expressed sequence Al115024
Mm.22149	succinate dehydrogenase complex,	25	Mm.199903	expressed sequence Al850290
	subunit A, flavoprotein (Fp)		Mm.199652	expressed sequence Al838505
Mm.2214	septin 4		Mm.198588	expressed sequence Al851970
Mm.220966	reticulon 4		Mm.19834	RIKEN cDNA 0610033L03 gene
Mm.220898	calmodulin 3		Mm.197523	brain acyl-CoA hydrolase
Mm.220885	neurochondrin	30	Mm.196614	eukaryotic translation elongation factor
Mm.2206	NADH dehydrogenase (ubiquinone) fla-			1 alpha 1
	voprotein 2		Mm.196611	synapsin I
Mm.219776	RIKEN cDNA 1110001E17 gene		Mm.196607	eukaryotic translation initiation factor 5A
Mm.218848	RIKEN cDNA 3010002G01 gene		Mm.196605	hexokinase 1
Mm.218764	guanine nucleotide binding protein 13,	35	Mm.196578	mitochondrial carrier homolog 1
	gamma		Mm.196344	lusterin
Mm.218611	receptor (calcitonin) activity modifying		Mm.196239	RIKEN cDNA 4922501H04 gene
M 04740	protein 2		Mm.195869	ATPase, H+ transporting, lysosomal
Mm.21743	malate dehydrogenase, mitochondrial			31kDa, V1 subunit E
Mm.216438	Mus musculus, clone IMAGE:5068657,	40	Mm.1956	neurofilament, light polypeptide
M 016040	mRNA, partial cds		Mm.19370	ATP synthase, H+ transporting, mi-
Mm.216240	Mus musculus, clone IMAGE:3594799,		M 400500	tochondrial F1F0 complex, subunit e
Mm 0140E	mRNA		Mm.193539	H1 histone family, member 2
Mm.21485 Mm.214549	RIKEN cDNA 2610102M01 gene	45	Mm.192991	Mus musculus, Similar to metallot-
WIIII.214549	Mus musculus, Similar to vesicle-asso-	45		hionein 1, clone MGC:27821 IMAGE:
	ciated calmodulin-binding protein, clone MGC:28873 IMAGE:4527857, mRNA,		Mm.19133	3483861, mRNA, complete cds amyloid beta (A4) precursor-like protein
	complete cds		WIIII. 19133	2
Mm.2133	centaurin, gamma 3		Mm.19047	expressed sequence Al425998
Mm.212672	S100 protein, beta polypeptide, neural	50	Mm.182912	growth hormone inducible transmem-
Mm.212516	RIKEN cDNA 2900002L20 gene	-	WIII. 102312	brane protein
	deleted in polyposis 1		Mm.18218	ganglioside-induced differentiation-as-
Mm.21251				sociated-protein 1
Mm.21251 Mm.21162	genes associated with retinoid-IFN-in-			
	genes associated with retinoid-IFN-in- duced mortality 19		Mm.181894	•
	genes associated with retinoid-IFN-in- duced mortality 19 transthyretin	55	Mm.181894 Mm.181721	RIKEN cDNA 2900092E17 gene
Mm.21162	duced mortality 19	55		RIKEN cDNA 2900092E17 gene RIKEN cDNA 2610041P16 gene
Mm.21162 Mm.2108	duced mortality 19 transthyretin	55	Mm.181721	RIKEN cDNA 2900092E17 gene
Mm.21162 Mm.2108 Mm.21071	duced mortality 19 transthyretin ADP-ribosylation-like 2	55	Mm.181721 Mm.180182	RIKEN cDNA 2900092E17 gene RIKEN cDNA 2610041P16 gene cytochrome c oxidase, subunit Vb

E	P 1	529	838	A1
---	-----	-----	-----	----

Mm.115124 brain protein 14 Mm.114810 expressed sequence AW060990 Mm.1147 Mus musculus calmodulin III (Calm3) mRNA, 3' untranslated region Mm.10727 ATPase, H+ transporting, lysosomal 56/58kD, V1 subunit B, isoform 2 Mm.103709 potassium inwardly-rectifying channel, subfamily J, member 10 Mm.103605 DnaJ (Hsp40) homolog, subfamily B, solute carrier family 25 (mitochondrial 10 member 10 Mm.102278 secretory carrier membrane protein 5 Mm.102244 expressed sequence R74975 Mm.101476 (Manual assignment) BNPI, VGLUT-1, mouse homolog of putative vesicular ESTs, Weakly similar to PBAS MOUSE 15 glutamate transporter, Na+/Phosphate cotransporter Mm.100980 calneuron 1 Mm.1008 prostaglandin D2 synthase (21 kDa.

18

Mm.157859 **ESTs** Mm.157648 RIKEN cDNA 5730403B10 gene Mm.15711 cyclic nucleotide phosphodiesterase 1 Mm.156959 beta-spectrin 4 Mm.15571 amyloid beta (A4) precursor protein Mm.15512 potassium voltage-gated channel, shaker-related subfamily, beta member 2 Mm.154651 purine rich element binding protein B Mm.153758 RIKEN cDNA 0610040H15 gene Mm.15125 stromal cell derived factor receptor 1

ribosomal protein S13

expressed sequence Al173355

RIKEN cDNA 2610009E16 gene

neurofilament, medium polypeptide

17

synuclein, alpha

tein A2/B1

tor), member 4

dynamin

creatine kinase, brain

aldolase 1, A isoform

Mus musculus, clone MGC:31632

IMAGE:4511454, mRNA, complete cds

heterogeneous nuclear ribonucleopro-

carrier; adenine nucleotide transloca-

RIKEN cDNA 2610301115 gene

RIKEN cDNA 5031406P05 gene

RIKEN cDNA 2410003L22 gene

PROBASIN PRECURSOR

Mm.177117

Mm.176927

Mm.17484

Mm.16831

Mm.16769

Mm.16767

Mm.16763

Mm.16228

Mm.16080

Mm.158871

Mm.157929

Mm.14798

Mm.142511

Mm.142187

Mm.142140

Mm.13859

Mm.12468

Mm.1239

Mm.1222

Mm.124592

Liste D: Positivmarker neurale Stammzellen (1.):

(Manual) Prostaglandin H2 D-Isomera-

se (PGD2 SYNTHASE) (PGDS2)

(Manual) copper transport protein/cha-

factor

receptor

(PGDS) member of lipocalin family

ES-Zellen +; PSA-NCAM - ; Adult brain -

perone ATOX1

[0042]

Mm.9703

Mm.1008

20

25

40

Mm.140761 DnaJ (Hsp40) homolog, subfamily C, member 5 Mm.139797 expressed sequence Al848587 Mm.139239 RIKEN cDNA 2900016C05 gene Mm.139239 RIKEN cDNA 2900016C05 gene Mm.139239 RIKEN cDNA 2900016C05 gene Mm.138866 apolipoprotein E

Mm.930 cathepsin L Mm.90787 nerve growth (TNFRSF16) associated protein 1 Mm.90587 enolase 1, alpha non-neuron Mm.90115 lysophospholipase 1 Mm.90003 gap junction membrane channel protein beta 3 Mm.88302 EST, Weakly similar to S14234 hypo-

Mm.1383 Rho GDP dissociation inhibitor (GDI) Mm.135621 expressed sequence Al848120 Mm.13445 3-oxoacid CoA transferase Mm.1339 chromogranin B Mm.131127 RIKEN cDNA 6230410L23 gene Mm.12958 kinesin light chain 2 Mm.12860 G protein-coupled receptor 37-like 1 Mm.1268 proteolipid protein (myelin) Mm.1268 (Manual assignment) PLP Myelin Proteolipid Protein, uh05d10.r1 Soares mouse hypothalamus NMHy Mus musculus cDNA clone 1617043 5' similar to gb:M54927 MYELIN PROTEOLIPID **PROTEIN**

glioblastoma amplified sequence

expressed sequence AW214631

brain abundant, membrane attached si-

glial fibrillary acidic protein

gnal protein 2

ribosomal protein L41

Mm.88212 tubulin, alpha 6 Mm.87581 (Manual) fibronectin 1, internal tag (major tag probably AAAAAAAAAA) Mm.87293 WD repeat domain 12

thetical protein - mouse

Mm.87216 Rab geranylgeranyl transferase, a sub-Mm.8155 TG interacting factor Mm.78861 nucleolar and coiled-body phosphoprotein 1

Mm.76780 **ESTs** Mm.7417 cyclin D3 RNA polymerase 1-4 (194 kDa subunit) Mm.7387 Mm.7381 hypoxia induced gene 1 Mm.725 ribosomal protein L7a **ESTs** Mm.71046

Mm.70127 ribosomal protein L12 Mm.69647 pancreas specific transcription factor,

1a

19

M 00040	-DNA A5455540			
Mm.69049	cDNA sequence AF155546			IMAGE:3992883, mRNA, complete cds
Mm.6700	eukaryotic translation initiation factor		Mm.3845	Mus musculus, eukaryotic translation
M 66	4E binding protein 1			termination factor 1, clone MGC:18745
Mm.66 Mm.6579	ribosomal protein S4, X-linked centromere autoantigen A	5	M 20151	IMAGE:3992883, mRNA, complete cds
Mm.6534	· ·	3	Mm.38151 Mm.38057	adenylosuccinate lyase
Mm.6343	calpain, small subunit 1			ESTs Man Alexander Alexander
Mm.584	nucleophosmin 1 annexin A2		Mm.3776	Mus musculus, clone MGC:37810
			M 0750	IMAGE:5098241, mRNA, complete cds
Mm.57223 Mm.57153	helicase, lymphoid specific	10	Mm.3752	RAN binding protein 1
Mm.5624	sterol O-acyltransferase 2	10		B-cell receptor-associated protein 37
WIIII.3024	DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 16		Mm.360 Mm.3572	cytochrome c oxidase, subunit Va
Mm.548	• • •		Mm.35621	RIKEN cDNA 1110033J19 gene
Mm.5305	cytochrome c oxidase, subunit Vic (Manual) GNB2L1, RACK1, Receptor		Mm.35605	ESTs
WIII.5505	of activated C kinase, WD40-repeat	15	Mm.3487	cadherin 1
	protein	,,,	Mm.3486	ribosomal protein L30 ribosomal protein L3
Mm.5290	(Manual) 60S ribosomal protein L17		Mm.34828	heat shock protein, 105 kDa
141111.0200	(L23) (popey3-annotation wrong)		Mm.34797	cellular retinoic acid binding protein I
Mm.4993	matrix metalloproteinase 3		Mm.34606	RIKEN cDNA 2610511F02 gene
Mm.493	CCCTC-binding factor	20	Mm.34554	Mus musculus, Similar to E2F trans-
Mm.4890	Finkel-Biskis-Reilly murine sarcoma vi-		14111.54554	cription factor 4, p107/p130-binding,
141111.4050	rus (FBR-MuSV) ubiquitously expres-			clone MGC:37558 IMAGE:4987691,
	sed (fox derived)			mRNA, complete cds
Mm.4770	frizzled homolog 7 (Drosophila)		Mm.3438	lamin A
Mm.4742	proliferation-associated 2G4, 38kD	25	Mm.34351	Mus musculus, Similar to hypothetical
Mm.46461	L-threonine dehydrogenase		141111.04001	protein FLJ13187, clone MGC:28979
Mm.4606	branched chain aminotransferase 1.			IMAGE:4503757, mRNA, complete cds
	cytosolic		Mm.34102	ornithine decarboxylase, structural
Mm.4560	low density lipoprotein receptor-related		Mm.3379	serine hydroxymethyl transferase 1
	protein associated protein 1	30		(soluble)
Mm.45237	RIKEN cDNA 2610318N02 gene		Mm.33240	epithelial V-like antigen
Mm.45151	RIKEN cDNA 1700043E15 gene		Mm.33202	RIKEN cDNA 2410018A17 gene
Mm.4502	mini chromosome maintenance de-		Mm.32879	testis expressed gene 17
	ficient (S. cerevisiae)		Mm.321	secreted phosphoprotein 1
Mm.43831	lectin, galactose binding, soluble 1	35	Mm.318	RIKEN cDNA 2010107E04 gene
Mm.43162	RIKEN cDNA 0710008D09 gene		Mm.31227	expressed sequence AW123847
Mm.42960	RIKEN cDNA 2610301D06 gene		Mm.30929	peroxiredoxin 1
Mm.4280	RIKEN cDNA 2010203J19 gene		Mm.3049	CDC28 protein kinase 1
Mm.42790	ribosomal protein S18		Mm.30242	peptidylprolyl isomerase D (cyclophilin
Mm.42767	ribosomal protein S17	40		D)
Mm.42197	proteasome (prosome, macropain)		Mm.30184	RIKEN cDNA 2700086123 gene
	subunit, beta type 1		Mm.30114	amyotrophic lateral sclerosis 2 (juveni-
Mm.42196	nuclear protein 95			le) homolog (human)
Mm.42195	RuvB-like protein 1		Mm.30060	RIKEN cDNA 2310008N12 gene
Mm.41467	Mus musculus, clone MGC:28892	45	Mm.30049	complement component 1, q subcom-
	IMAGE:4912251, mRNA, complete cds			ponent binding protein
Mm.41151	ESTs		Mm.30034	translocase of inner mitochondrial
Mm.41061	RIKEN cDNA 1810055P05 gene			membrane 8 homolog a (yeast)
Mm.41	(Manual) Mitochondrial ATP synthase		Mm.29904	mitochondrial ribosomal protein L15
	oligomycin sensitivity conferral protein	50	Mm.29902	Mus musculus, Similar to phosphoseri-
	(OSCP) (ATP5O)			ne aminotransferase, clone MGC:6462
Mm.4095	inactive X specific transcripts			IMAGE:2616298, mRNA, complete cds
Mm.4024	cofilin 1, non-muscle		Mm.29859	eukaryotic translation initiation factor 2,
Mm.3925	S100 calcium binding protein A4			subunit 2 (beta, 38kDa)
Mm.38718	ESTs, Moderately similar to S12207 hy-	55	Mm.29856	RIKEN cDNA 9130022B02 gene
	pothetical protein		Mm.29717	3-monooxgenase/tryptophan 5-mo-
Mm.3845	Mus musculus, eukaryotic translation			nooxgenase activation protein, gamma
	termination factor 1, clone MGC: 18745			polypeptide

21

Mm 20714	(Manual) mayor version of mysels			4-2
Mm.29714	(Manual) mouse version of muscle- specific protein M9		M 04506	tor), member 13
Mm.29675	thioredoxin-like 2		Mm.24506	Mus musculus, clone IMAGE:3591061,
Mm.29619	RIKEN cDNA 1200007E24 gene		Mm 2427	mRNA, partial cds
Mm.29513	•	5	Mm.2437 Mm.2424	BING4 protein
WIIII.29515	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 7 (14.5kD, B14.5a)	3	•	ribosomal protein L10A
Mm.29504			Mm.24220	RIKEN cDNA 2310003F16 gene
Mm.2942	sperm specific antigen 1		Mm.24219	RIKEN cDNA 1810037117 gene
Mm.29405	asparagine synthetase		Mm.24174	Mus musculus, similar to alanyi-tRNA
Mm.29363	ring-box 1 RIKEN cDNA 2310044F10 gene	10		synthetase (H. sapiens), clone MGC:
Mm.2930	•	10		37368 IMAGE:4976684, mRNA, com-
WIII1.2950	Mus musculus, Similar to peter pan		Mm 000E	plete cds
	(Drosophila) homolog, clone MGC: 25669 IMAGE:4489442, mRNA, com-		Mm.2395	male enhanced antigen 1
	plete cds		Mm.2355	prohibitin
Mm.29192	asparaginyl-tRNA synthetase	15	Mm.235	ubiquitin B
Mm.29148	RIKEN cDNA 2400008B06 gene	15		integrin beta 4 binding protein
Mm.29122	RIKEN cDNA 0610012D09 gene		Mm.22626	Mus musculus, Similar to chromosome
Mm.29076	RIKEN cDNA 0010012D09 gene			14 open reading frame 3, clone MGC:
Mm.28919	destrin			36589 IMAGE:5320590, mRNA, com-
Mm.28892	expressed sequence AA959742	20	Mm.2246	plete cds proteasome (prosome, macropain)
Mm.28805	SET translocation	20	WIII1.2240	
Mm.2849	heat shock protein, 74 kDa, A		Mm.22421	subunit, beta type 7 telomerase binding protein, p23
Mm.28483	Mus musculus, Similar to hypothetical		Mm.22421	telomerase binding protein, p23
1011111.20-400	protein FLJ22479, clone IMAGE:		Mm.22317	RIKEN cDNA 8430410A17 gene
	4487274, mRNA, partial cds	25		cyclin D1
Mm.28405	serum/glucocorticoid regulated kinase		Mm.22271	smt3-specific isopeptidase 1
Mm.28173	ESTs, Moderately similar to JC5224		Mm.220992	Mus musculus, clone IMAGE:3492506,
	methioninetRNA ligase		WIII.220332	mRNA, partial cds
Mm.28053	RIKEN cDNA 1110017C15 gene		Mm.219671	Mus musculus, clone MGC:36369
Mm.28035	ESTs, Weakly similar to	30		IMAGE:4982239, mRNA, complete cds
	TRHY_HUMAN TRICHOHYALI		Mm.219458	RNA binding protein gene with multiple
Mm.27901	RIKEN cDNA 1110020J08 gene			splicing
Mm.27858	RIKEN cDNA 1110036B12 gene		Mm.218533	RIKEN cDNA 1500016H10 gene
Mm.27855	replication factor C (activator 1) 2		Mm.2180	heat shock protein, 84 kDa 1
	(40kD)	35	Mm.21758	cytochrome P450, 2e1, ethanol induci-
Mm.2758	makorin, ring finger protein, 3			ble
Mm.27536	ESTs, Highly similar to hypothetical		Mm.21630	expressed sequence AU022237
	protein FLJ14075		Mm.21569	RIKEN cDNA 2700069E09 gene
Mm.27526	(Manual) Arginyl tRNA synthetase (RI-		Mm.213020	(Manual) 60S ribosomal protein L32
	KEN cDNA 2610011N19)	40		(RPL32)
Mm.27186	Mus musculus, Similar to CG7083 ge-		Mm.212899	Mus musculus, Similar to RIKEN cDNA
	ne product, clone MGC:6480 IMAGE:			1200009K13 gene, clone MGC: 18794
	2646515, mRNA, complete cds			IMAGE:4193513, mRNA, complete cds
Mm.2718	eukaryotic translation elongation factor		Mm.21289	ribosomal protein S12
	1 beta 2	45	Mm.21086	eukaryotic translation elongation factor
Mm.2718	eukaryotic translation elongation factor			1 delta (guanine nucleotide exchange
	1 beta 2			protein)
Mm.27134	RIKEN cDNA 2610033C09 gene		Mm.210638	EST
Mm.265	ribosomal protein S25		Mm.21062	expressed sequence C87860
Mm.2647	profilin 1	50	Mm.21054	nuclease sensitive element binding
Mm.2623	serine (or cysteine) proteinase inhibitor,			protein 1
	clade B (ovalbumin), member 6		Mm.20943	FK506 binding protein 9
Mm.25642	RIKEN cDNA 2310034K10 gene		Mm.20925	G1 to phase transition 1
Mm.254	tumor protein, translationally-controlled		Mm.20918	nuclear localization signal protein ab-
	1	55		sent in velo-cardio-facial patients
Mm.25328	ESTs		Mm.20848	regulatory factor X-associated ankyrin-
Mm.24513	solute carrier family 25 (mitochondrial			containing protein
	carrier; adenine nucleotide transloca-		Mm.20847	sorting nexin 5

EP 1 529 838 A1

Mm.20294	colonarhambata ayathataa 0		M 457770	DUCEN BALL COLORS (E.C.
	selenophosphate synthetase 2		Mm.157778	RIKEN cDNA 2610034E13 gene
Mm.20290	expressed sequence AW536573		Mm.154915	ribosomal protein S29
Mm.20288	glutathione reductase 1		Mm.154387	transketolase
Mm.200920	ribosomal protein S28		Mm.153963	CD8 antigen, beta chain
Mm.197601	heat shock 10 kDa protein 1 (chapero-	5	Mm.153159	chaperonin subunit 6a (zeta)
	nin 10)		Mm.152291	EST
Mm.197555	hypothetical protein MGC6664		Mm.151329	karyopherin (importin) beta 3
Mm. 197551	heat shock 70kD protein 8		Mm.148973	RIKEN cDNA 3010025E17 gene
Mm.196604	angio-associated migratory protein, re-		Mm.147946	MYB binding protein (P160) 1a
	lated sequence	10	Mm.147693	ribosomal protein S3
Mm.196586	cullin 2		Mm.14768	reduced expression 3
Mm. 196581	mitogen activated protein kinase 1		Mm.14663	ATP synthase, H+ transporting, mi-
Mm.196526	ADP-ribosylation factor 6			tochondrial F0 complex, subunit g
Mm.196396	tubulin, alpha 1		Mm.143141	eukaryotic translation initiation factor
Mm.196081	peptidylprolyl isomerase (cyclophilin)-	15		1A
	like 1		Mm.142740	metallothionein 2
Mm.196	neural precursor cell expressed, de-		Mm.14245	ribosomal protein, large P2
	velopmentally down-regulated gene 8		Mm.14244	ribosomal protein L9
Mm.195894	Mus musculus, clone MGC:11792			
WIIII. 133034		20	Mm.141443	lactate dehydrogenase 1, A chain
Mm.19169	IMAGE:3595167, mRNA, complete cds	20	Mm.141187	trans-golgi network protein 2
	thioredoxin-like (32kD)		Mm.140380	ribosomal protein L23
Mm.188	(Manual) X-linked phosphoglycerate ki-		Mm.139825	Mus musculus, Similar to xylosylprotein
11	nase (PGK1)			betal,4-galactosyltransferase, poly-
Mm.18637	teratocarcinoma expressed, serine rich			peptide 7 (galactosyltransferase I), clo-
Mm. 18459	fibroblast growth factor inducible 14	25		ne MGC: 28643 IMAGE:4224150, mR-
Mm.183022	DNA segment, Chr 8, Brigham & Wo-			NA, complete cds
	men's Genetics 1112 expressed		Mm.13705	(Manual) mouse version of p180 ribo-
Mm.182951	proteasome (prosome, macropain)			some receptor/ribosome binding prote-
	subunit, alpha type 2			in 1 RRBP1
Mm.182931	phosphoribosylaminoimidazole car-	30	Mm.13020	ribosomal protein L13a
	boxylase, phosphoribosylaminoribosy-		Mm.12909	amyloid beta (A4) precursor protein-
	laminoimidazole, succinocarboxamide			binding, family A, member 3
	synthetase		Mm.1275	thioredoxin 1
Mm.182471	RIKEN cDNA 2610524G07 gene		Mm.12508	karyopherin (importin) alpha 2
Mm.181765	Mus musculus 8 days embryo whole	35	Mm.1164	SEC61, gamma subunit (S. cerevisiae)
	body cDNA, RIKEN full-length enriched		Mm.11376	ribosomal protein L36
	library, clone:5730409M10:CCAAT/en-		Mm.1125	expressed in non-metastatic cells 2,
	hancer binding protein alpha (C/EBP),			protein (NM23B) (nucleoside diphos-
	related sequence 1, full insert se-			phate kinase)
	quence	40	Mm.1120	endometrial bleeding associated factor
Mm.181740	interferon-related developmental regu-		Mm.108076	phosphofructokinase, platelet
	lator 2		Mm.10706	RIKEN cDNA 2010004J23 gene
Mm.180299	DNA segment, Chr 16, Wayne State		Mm.10706	
141111.100255	University 109, expressed		WIIII. 10708	(Manual) mouse version of 60S riboso-
Mm.17932	purine-nucleoside phosphorylase	45	Mm 10700	mal protein L4
Mm.1777		43	Mm.10702	calcyclin binding protein
Mm.176845	heat shock protein, 60 kDa		Mm.10665	Mus musculus, cione IMAGE:3498496,
	RIKEN cDNA 1110069M14 gene			mRNA, partial cds
Mm.175848	(Manual) small Ca-binding protein Cal-		Mm.10623	expressed sequence Al480570
	gizzarin (S100A11) (ENDOTHELIAL		Mm.10600	glutamate dehydrogenase
	MONOCYTE-ACTIVATING POLYPEP-	50	Mm.1056	solute carrier family 1, member 7
M 475004	TIDE) (EMAP)		Mm.10474	RIKEN cDNA 3110005M08 gene
Mm.175661	RIKEN cDNA 1110036C17 gene		Mm.101619	EST
Mm.1710	hydroxymethylbilane synthase		Mm.10	spermidine synthase
Mm.17031	POU domain, class 5, transcription fac-		Mm.4325	Kruppel-like factor 4 (gut) [Swissprot:
	tor 1	55		splQ60793;splQ9R255;]
Mm.16757	solute carrier family 20, member 1		Mm.12919	insulin-like growth factor 2, binding pro-
Mm.1639	myeloid cell leukemia sequence 1			tein 1 [Swissprot: splO88477;]
Mm.16110	cyclin E1		Mm.20348	nidogen 2 [Swissprot: splO88322;

EP 1	529	838 /	11
			••

	aniORREGO:aniOOCTO4:1		Mm 7700	
Mm.34407	splQ8R5G0;splQ9CT94;] MAD homolog 7 (Drosophila) [Swiss-		Mm.7793	protein phosphatase 1, catalytic subu-
101111.54407	prot: splO35253;splQ9CSC7;]		Mm.7723	nit, gamma isoform poly(A) binding protein, nuclear 1
Mm.4451	hairy and enhancer of split 1, (Droso-		Mm.76278	RIKEN cDNA 2610203K23 gene
	phila) [Swissprot: none]	5	Mm.7516	nuclear autoantigenic sperm protein (hi-
Mm.57195	nodal [Swissprot: splP43021;]			stone-binding)
Mm.1249	laminin, gamma 1 [Swissprot: spl		Mm.7312	DNA segment, Chr 17, human D6S56E
	P02468;]			2
Mm.27706	ash2 (absent, small, or homeotic)-like		Mm.7141	proliferating cell nuclear antigen
	(Drosophila) [Swissprot:	10	Mm.6787	splicing factor, arginine/serine-rich 3
Mm.4603	scavenger receptor class B1 [Swiss-			(SRp20)
	prot: splQ61009;splQ9CWJ7;}		Mm.66	ribosomal protein S4, X-linked
Mm.181562	adhesion regulating molecule 1 [Swis-		Mm.6476	RIKEN cDNA 2700084L22 gene
	sprot: splQ8VCl8;splQ922A7;		Mm.64104	RIKEN cDNA 2410016F19 gene
	splQ9JKV1;]	15	Mm.6343	nucleophosmin 1
Mm.43444	MAD2 (mitotic arrest deficient, homo-		Mm.61901	expressed sequence Al429604
14 400075	log)-like 1 (yeast) [Swissprot:		Mm.6065	inosine 5'-phosphate dehydrogenase 2
Mm.103675	sacsin [Swissprot: none]		Mm.5624	DEAD/H (Asp-Glu-Ala-Asp/His) box po-
Mm.980	tenascin C [Swissprot: splQ64706;			lypeptide 16
M 5000	splQ9WUU4;]	20	Mm.548	cytochrome c oxidase, subunit VIc
Mm.5090	cripto, teratocarcinoma-derived growth		Mm.5305	guanine nucleotide binding protein, beta
Mm.30177	factor (Tdgf1) D11Ertd603e, DNA segment, Chr 11,		Mm FOF	2, related sequence 1
WIIII.30177	ERATO Doi 603		Mm.525	eukaryotic translation initiation factor 4, gamma 2
Mm.233844	C330012H03Rik, RIKEN cDNA	25	Mm.5114	dishevelled 2, dsh homolog (Drosophi-
WWW.2000-1-7	C330012H03		WIII.5114	la)
	000012.700		Mm.4933	mini chromosome maintenance de-
Liste E: Posit	ivmarker neurale Stammzellen (2.);			ficient 6 (S. cerevisiae)
	· ''		Mm.4890	•
			WII 11.4030	rinker-diskis-Helliv murine sarcoma vi-
ES-Zellen +; l	PSA-NCAM -/+; Adult brain -	30	WIII.4050	Finkel-Biskis-Reilly murine sarcoma virus (FBR-MuSV) ubiquitously expres-
ES-Zellen +; I	PSA-NCAM -/+; Adult brain -	30	WIII.4030	•
ES-Zellen +; [0043]	PSA-NCAM -/+; Adult brain -	30	Mm.4846	rus (FBR-MuSV) ubiquitously expres-
[0043]		30		rus (FBR-MuSV) ubiquitously expressed (fox derived)
[0043] Mm.99776	cytosolic aminopeptidase P	30	Mm.4846	rus (FBR-MuSV) ubiquitously expres- sed (fox derived) lamin B1
[0043] Mm.99776 Mm.9916	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene	<i>30</i>	Mm.4846 Mm.4756 Mm.46754 Mm.46533	rus (FBR-MuSV) ubiquitously expres- sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene
[0043] Mm.99776 Mm.9916 Mm.99	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2		Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551	rus (FBR-MuSV) ubiquitously expres- sed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene		Mm.4846 Mm.4756 Mm.46754 Mm.46533	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1
[0043] Mm.99776 Mm.9916 Mm.99	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre-		Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y310, similar	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family		Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re-	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD)	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.45132	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin-	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homo-
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast)
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin-	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.45132	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro-	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast)
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa	35	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767 Mm.4237	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6	35 40 45	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767 Mm.4237	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212 Mm.880	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6	35 40 45	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.4237 Mm.42197	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212 Mm.880 Mm.8552	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5	35 40 45	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.4237 Mm.42197	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212 Mm.880	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, si-	35 40 45	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.4237 Mm.42197 Mm.4215 Mm.41940 Mm.4189 Mm.41023	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.88136 Mm.88212 Mm.880 Mm.8552 Mm.8256	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, si- gnal transduction associated 1	35 40 45	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.4237 Mm.42197 Mm.4215 Mm.4215 Mm.41940 Mm.4189	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibo-
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.88212 Mm.880 Mm.8552 Mm.8256 Mm.8155	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, si- gnal transduction associated 1 TG interacting factor	35 40 45	Mm.4846 Mm.4756 Mm.46754 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.4237 Mm.42197 Mm.4215 Mm.41940 Mm.4189 Mm.41023 Mm.41023 Mm.4078	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene viillin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibody Ki 67
[0043] Mm.99776 Mm.9916 Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.88136 Mm.88212 Mm.880 Mm.8552 Mm.8256	cytosolic aminopeptidase P RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre- sponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re- gulated protein, 78kD) high mobility group nucleosomal bin- ding domain 2 heterogeneous nuclear ribonucleopro- tein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, si- gnal transduction associated 1	35 40 45	Mm.4846 Mm.4756 Mm.46754 Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45132 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.4237 Mm.42197 Mm.4215 Mm.41940 Mm.4189 Mm.41023	rus (FBR-MuSV) ubiquitously expressed (fox derived) lamin B1 leptin receptor expressed sequence Al316867 RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibo-

EP	1	529	838	A
	EP	EP 1	EP 1 529	EP 1 529 838

M 4004	****			
Mm.4024	cofilin 1, non-muscle		Mm.29122	RIKEN cDNA 0610012D09 gene
Mm.3931	Max protein		Mm.29055	chromobox homolog 1 (Drosophila HP1
Mm.38930	expressed sequence AA407558			beta)
Mm.38912	RIKEN cDNA 2410129H14 gene		Mm.29054	RIKEN cDNA 2610529112 gene
Mm.38611	RIKEN cDNA 2210021A15 gene	5	Mm.29005	expressed sequence AU021749
Mm.38528	RIKEN cDNA 2810430M08 gene		Mm.28995	RIKEN cDNA 2010009J12 gene
Mm.38306	macrophage erythroblast attacher		Mm.28985	ribosomal protein L27
Mm.3797	nucleosome assembly protein 1-like 1		Mm.28965	RIKEN cDNA 0710007A14 gene
Mm.37835	ribosomal protein L7		Mm.28964	Mus musculus, clone IMAGE:4949762,
Mm.372	ribosomal protein S26	10		mRNA, partial cds
Mm.36511	mitochondrial ribosomal protein L32		Mm.28961	cleavage and polyadenylation specific
Mm.35844	growth arrest specific 5			factor 5, 25 kD subunit
Mm.35829	erythroid differentiation regulator		Mm.28909	protein tyrosine phosphatase 4a1
Mm.35661	Mus musculus, Similar to hypothetical		Mm.28899	RIKEN cDNA 1110059P08 gene
	protein, clone MGC:29235 IMAGE:	15		SET translocation
	5043282, mRNA, complete cds		Mm.28805	SET translocation
Mm.35087	expressed sequence AA673488		Mm.28805	SET translocation
Mm.3501	kinesin family member C5A		Mm.28726	
Mm.34914	ESTs			EST C77032
Mm.3487	ribosomal protein L30	20	Mm.28694	RIKEN cDNA 2410088K19 gene
Mm.3444	bromodomain-containing 2	20	Mm.28560	Ly1 antibody reactive clone
Mm.34385			Mm.28499	Mus musculus, similar to CG15881 ge-
	expressed sequence Al450344			ne product (H. sapiens), clone MGC:
Mm.34261	expressed sequence AW557761			36308 IMAGE:5040108, mRNA, com-
Mm.3381	ribosomal protein S8			plete cds
Mm.3380	kinesin family member 5B	25	Mm.28299	ESTs, Highly similar to GUAA_HUMAN
Mm.3360	tyrosine 3-monooxygenase/tryptophan			GMP SYNTHASE
	5-monooxygenase activation protein,		Mm.28222	RIKEN cDNA 2610307C23 gene
	zeta polypeptide		Mm.28121	RIKEN cDNA 1110061A19 gene
Mm.326	RIKEN cDNA 1110038L14 gene		Mm.28044	filamin-like protein
Mm.320	DNA polymerase alpha 2, 68 kDa	30	Mm.27972	NS1-associated protein 1
Mm.3199	RIKEN cDNA 1500001N04 gene		Mm.27927	heterogeneous nuclear ribonucleopro-
Mm.31512	ring finger protein 2			tein A1
Mm.31228	RIKEN cDNA 1810022K09 gene		Mm.27852	expressed sequence AW555814
Mm.30806	ribosomal protein L19		Mm.27818	eukaryotic translation elongation factor
Mm.3054	alpha-L-iduronidase	35		2
Mm.3035	RIKEN cDNA 3110006P09 gene		Mm.27796	RIKEN cDNA 5730427N09 gene
Mm.30270	proteasome (prosome, macropain) sub-		Mm.27669	small nuclear ribonucleoprotein E
	unit, alpha type 4		Mm.27660	RIKEN cDNA 5730420G12 gene
Mm.30120	ribosomal protein S27-like		Mm.27624	RIKEN cDNA C530002L11 gene
Mm.30069	RIKEN cDNA 1200003J11 gene	40	Mm.27293	RIKEN cDNA 4833420K19 gene
Mm.30011	ribosomal protein S23		Mm.27269	RIKEN cDNA 2310037I24 gene
Mm.29931	cell division cycle 20 homolog (S. cere-		Mm.27141	Rac GTPase-activating protein 1
	visiae)		Mm.27074	RIKEN cDNA 2610019N13 gene
Mm.29923	SMT3 (supressor of mif two, 3) homolog		Mm.265	ribosomal protein S25
	2 (S. cerevisiae)	45	Mm.2591	RNA binding motif protein 3
Mm.29911	RIKEN cDNA 3200001M24 gene		Mm.25558	RIKEN cDNA 2410018J24 gene
Mm.29896	ribosomal protein L21		Mm.25542	(Manual) strange EST contig in intron of
Mm.2986	expressed sequence AW146116			p137 (GPI-anchored transcytosis prote-
Mm.29829	expressed sequence Al326010			in), maybe alternative C-terminus of
Mm.29666	solute carrier family 25 (mitochondrial	50		splQ60865
	carnitine/acylcarnitine translocase),		Mm.254	tumor protein, translationally-controlled
	member 20 .		***************************************	1
Mm.2966	isocitrate dehydrogenase 2 (NADP+),		Mm.25299	ESTs, Weakly similar to simple repeat
···	mitochondrial			sequence-containing transcript
Mm.29296	RIKEN cDNA 1110003H02 gene	55	Mm.25164	gene trap locus 1-13
Mm.29194	RIKEN cDNA 1700094M07 gene		Mm.25137	RIKEN cDNA 2410004B18 gene
Mm.29133	budding uninhibited by benzimidazoles		Mm.24870	(Manual assignment) UBP7 ubiquitin
20100	1 homolog, beta (S. cerevisiae)		141111.240/U	hydrolase
	i nomolog, beta (o. celevisiae)			nyurolase

M 04504	A1115 40070			
Mm.24591	expressed sequence AW546279		Mm.21054	nuclease sensitive element binding pro-
Mm.2424	ribosomal protein L10A			tein 1
Mm.24219	RIKEN cDNA 1810037I17 gene		Mm.20927	transforming growth factor beta 1 indu-
Mm.24042	RIKEN cDNA 1210001E11 gene	_		ced transcript 4
Mm.23943	vesicle-associated membrane protein,	5	Mm.206399	ESTs
M 00750	associated protein A (33 kDa)		Mm.2038	Ras-GTPase-activating protein SH3-
Mm.23758	RIKEN cDNA 1110008P04 gene			domain binding protein
Mm.23695	dihydrofolate reductase		Mm.2025	survival motor neuron
Mm.23692	casein kinase II, alpha 1 related se-		Mm.200837	Mus musculus, clone IMAGE:5355658,
14 00000	quence 4	10		mRNA
Mm.23096	protein phosphatase 2 (formerly 2A), re-		Mm.196614	eukaryotic translation elongation factor
Mm 0007	gulatory subunit B", alpha		14 400000	1 alpha 1
Mm.2287	proteasome (prosome, macropain) sub-		Mm.196608	expressed sequence AA407306
Mm 00701	unit, alpha type 5	45	Mm.196526	ADP-ribosylation factor 6
Mm.22731	integrin beta 4 binding protein	15	Mm.196515	DNA segment, Chr 1, ERATO Doi 692,
Mm.2265 Mm.22387	U1 small nuclear ribonucleoprotein 1C		N 100000	expressed
	expressed sequence Al314668		Mm.196396	tubulin, alpha 1
Mm.22269	exportin 1, CRM1 homolog (yeast)		Mm.196365	RIKEN cDNA 4833416109 gene
Mm.22214	RIKEN cDNA 2610008F03 gene	20	Mm.196328	RIKEN cDNA 5830466J11 gene
Mm.220918	heterogeneous nuclear ribonucleopro- tein D-like	20	Mm.195898	phosphatidylethanolamine binding pro-
Mm.220342	Mus musculus, clone IMAGE:3669867.		Mm 1051	tein
WIIII.220342	mRNA, partial cds		Mm.1951	ribonucleic acid binding protein S1
Mm.219670	Mus musculus, Similar to eukaryotic		Mm.1948	t-complex testis expressed 1
WIII.213070	translation initiation factor 4 gamma, 1,	25	Mm.193688	RIKEN cDNA 2700059D21 gene
	clone IMAGE:4950789, mRNA, partial	25	Mm.19187	prothymosin alpha
	cds		Mm.19101	DEAD (aspartate-glutamate-alanine-
Mm.219668	RIKEN cDNA 2610209F03 gene		Mm.19015	aspartate) box polypeptide 5 serine racemase
Mm.219648	Mus musculus, Similar to nuclear matrix		Mm.18923	
141111.213040	protein p84, clone MGC:28284 IMAGE:	30	WITT. 10323	mini chromosome maintenance de- ficient 7 (S. cerevisiae)
	4010605, mRNA, complete cds	00	Mm.18921	valosin containing protein
Mm.21964	Mus musculus, clone IMAGE:3485208,		Mm. 18856	mitogen-activated protein kinase 6
101111111111111111111111111111111111111	mRNA, partial cds		Mm.18705	vacuolar protein sorting 4b (yeast)
Mm.21873	retroviral integration site 1		Mm.18700	RIKEN cDNA 1200009K13 gene
Mm.218657	cerebellar ataxia 3	35	Mm.18637	teratocarcinoma expressed, serine rich
Mm.21841	splicing factor, arginine/serine-rich 2		Mm.18516	H3 histone, family 3B
	(SC-35)		Mm.1843	heat shock protein, 86 kDa 1
Mm.218240	Mus musculus, clone IMAGE:5342828,		Mm.183102	actin-related protein 3 homolog (yeast)
	mRNA, partial cds		Mm.183016	thymine DNA glycosylase
Mm.2180	heat shock protein, 84 kDa 1	40	Mm.181880	RIKEN cDNA 1110007A14 gene
Mm.21764	small nuclear ribonucleoprotein poly-		Mm.181562	adhesion regulating molecule 1
	peptide G		Mm.1815	cytidine 5'-triphosphate synthase
Mm.21714	RIKEN cDNA 2410003A14 gene		Mm.180873	RIKEN cDNA 2510019J09 gene
Mm.21559	non-POU-domain-containing, octamer		Mm.180873	(Manual) probably reverse tag of 60S ri-
	binding protein	45		bosomal protein L18a
Mm.213452	Mus musculus, clone IMAGE:5320271,		Mm.180409	ubiquitin-conjugating enzyme E2H
	mRNA, partial cds		Mm.180271	RIKEN cDNA 5630400D24 gene
Mm.213020	(Manual) 60S ribosomal protein L32		Mm.17989	chaperonin subunit 8 (theta)
	(RPL32)		Mm.1777	heat shock protein, 60 kDa
Mm.21295	expressed sequence AW214031	50	Mm.1775	hematological and neurological expres-
Mm.21289	ribosomal protein S12			sed sequence 1
Mm.21281	ring finger protein 4		Mm.177451	RIKEN cDNA 5730544L10 gene
Mm.21185	adaptor-related protein complex AP-3,		Mm.17330	ESTs
	beta 1 subunit		Mm.17306	tropomyosin 3, gamma
Mm.2115	heterogeneous nuclear ribonucleopro-	55	Mm.1703	tubulin, beta 5
	tein U		Mm.16976	TAF9 RNA polymerase II, TATA box bin-
Mm.21094	DNA segment, Chr 9, Wayne State Uni-			ding protein (TBP)-associated factor, 32
	versity 138, expressed			kDa

31

Mm.16775 ribosomal protein S24 Mm.16767 heterogeneous nuclear ribonucleoprotein A2/B1 Mm.16711 mini chromosome maintenance de-5 ficient 2 (S. cerevisiae) Mm.16525 polo-like kinase homolog, (Drosophila) Mm.1639 myeloid cell leukemia sequence 1 Mm.16323 eukaryotic translation initiation factor 4A2 Mm.16323 eukaryotic translation initiation factor 10 4A2 Mm.156892 heterogeneous nuclear ribonucleoprotein D Mm.15571 amyloid beta (A4) precursor protein Mm.154915 ribosomal protein S29 15 Mm.153457 RIKEN cDNA 2810406C15 gene Mm.148973 RIKEN cDNA 3010025E17 gene Mm.142872 heterogeneous nuclear ribonucleoprotein K Mm.14245 20 ribosomal protein, large P2 Mm.14244 ribosomal protein L9 Mm.142363 RIKEN cDNA 2810036L13 gene Mm.140804 Mus musculus, guanine nucleotide binding protein (G protein), gamma 5, clone MGC:8292 IMAGE:3593324, mRNA, 25 complete cds Mm.140380 ribosomal protein L23 Mm.13886 suppressor of initiator codon mutations, related sequence 1 (S. cerevisiae) Mm.133825 RIKEN cDNA 0610010123 gene 30 Mm.13356 RIKEN cDNA 3110079L04 gene Mm.131705 Mus musculus, Similar to single-stranded DNA binding protein, clone MGC: 41439 IMAGE: 1314987, mRNA, complete cds 35 Mm.12858 eukaryotic translation initiation factor 4A1 Mm.12706 Mus musculus, Similar to CG11246 gene product, clone MGC:8248 IMAGE: 3591968, mRNA, complete cds 40 Mm.12604 sirtuin 1 ((silent mating type information regulation 2, homolog) 1 (S. cerevisiae) Mm.12568 expressed sequence AW541137 Mm.12508 karyopherin (importin) alpha 2 Mm.12441 expressed sequence AU014645 45 Mm.124 thymopoietin Mm.12236 zinc finger protein 207 Mm.12145 retinoblastoma binding protein 4 Mm.116989 actin-like Mm 111 poly(rC) binding protein 2 50 Mm.10706 RIKEN cDNA 2010004J23 gene Mm.10474 RIKEN cDNA 3110005M08 gene Mm.10409 golgi autoantigen, golgin subfamily a, 4 Mm.103675 sacsin Mm.1013 ligase I, DNA, ATP-dependent 55 Mm.101274 RIKEN cDNA 2010008E23 gene Mm.10076 mitochondrial ribosomal protein L13

Nmycl, neuroblastoma myc-related on-

Mm.16469

cogene 1

Patentansprüche

 Zellpopulation, dadurch gekennzelchnet, dass mindestens 5% der Zellen neurale Vorläuferzellen sind, die wenigstens einen der in Liste A oder Liste B aufgeführten Marker aufweisen.

32

- Zellpopulation, dadurch gekennzelchnet, dass mindestens 5% der Zellen, neurale Vorläuferzellen sind, die wenigstens zwei, bevorzugt wenigstens 3 der in Liste A oder Liste B aufgeführten Marker aufweisen.
- Zellpopulation, nach mindestens einem der Ansprüche 1 bis 2, dadurch gekennzelchnet, dass die neuralen Vorläuferzellen keinen in Liste C aufgeführten Marker aufweisen.
- Zellpopulation nach mindestens einem der Ansprüche 1 bis 3, dadurch gekennzeichnet, dass mindestens 25 % der Zellen neurale Vorläuferzellen sind.
- Zellpopulation nach mindestes einem der Ansprüche 1 bis 4, dadurch gekennzeichnet, dass es sich um eine murine Zellpopulation handelt und/oder die neuralen Vorläuferzellen aus Hirngewebe erhältlich ist.
- Verfahren zur Isolierung einer Zellpopulation nach mindestens einem der Ansprüche 1 bis 5 mit folgenden Schritten:
 - a) Entnahme einer Probe aus dem Hirn
 - b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von embryonalen Stammzellen zu neuralen Vorläuferzellen,
- b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- a) Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Vorläuferzellen,
- b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

a) Differenzierung von adulten, neuralen Stammzellen zu neuralen Vorläuferzellen,

33

b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von immortalisierten Zellen zu neuralen Vorläuferzellen.
- b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker.
- Verwendung mindestens eines Markers ausgewählt aus der Liste A oder Liste B zu Identifizierung oder Isolierung von neuralen Vorläuferzellen.
- Antikörper gegen einen Marker aus der Liste A, B oder C.
- Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste A, B oder C.
- Arzneimittel enthaltend die Zellpopulation nach einem der Ansprüche 1 bis 5.
- Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens einen der in Liste D oder Liste E aufgeführten Marker aufweisen.
- Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens zwei, bevorzugt wenigstens 3 der in Liste D oder Liste E aufgeführten Marker aufweisen.
- 13. Zellpopulation, nach mindestens einem der Ansprüche 11 bis 12, dadurch gekennzeichnet, dass die neuralen Stammzellen keinen in Liste A oder Liste C aufgeführten Marker aufweisen.
- Zellpopulation nach mindestens einem der Ansprüche 11-13, dadurch gekennzelchnet, dass mindestens 25% der Zellen neurale Stammzellen sind.
- 15. Zellpopulation nach mindestes einem der Ansprüche 11 bis 14, dadurch gekennzeichnet, dass es sich um eine murine Zellpopulation handelt und/oder die neuralen Stammzellen aus Hirngewebe erhältlich
- 16. Verfahren zur Isolierung einer Zellpopulation nach mindestens einem der Ansprüche 11 bis 15 mit folgenden Schritten:
 - a) Entnahme einer Probe aus dem Hirn

b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

34

oder

5

10

20

- a) Differenzierung von embryonalen Stammzellen zu neuralen Stammzellen,
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Stammzellen
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) De-Differenzierung von adulten, neuralen Vorläuferzeilen zu neuralen Stammzellen,
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von immortalisierten Zellen zu neuralen Stammzellen,
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker.
- Antikörper gegen einen Marker aus der Liste D, E, A oder C.
- 18. Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste D, E, A oder C.
- Arzneimittel enthaltend die Zellpopulation nach einem der Ansprüche 11 bis 15.



Europäisches Patentamt

Europäisches EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung

der nach Regel 45 des Europäischen Patemübereinkommens für das weitere Verfahren als europäischer Recherchenbericht gilt

EP 03 02 5506

Kategorie	Kennzeichnung des Dokur der maßgebliche	ments mit Angabe, soweit erforderlich	Betrifft Approach	KLASSIFIKATION DER ANMELDUNG (Int.CI.7)
х	ARSENIJEVIC YVAN Emultipotent neural the cortex of the a EXPERIMENTAL NEUROL Bd. 170, Nr. 1, July Seiten 48-62, XP002 ISSN: 0014-4886 * Seite 52, linke Srechte Spalte, Absarches	T AL: "Isolation of precursors residing in adult human brain" LOGY, li 2001 (2001-07), 2275728 Spalte, letzter Absatz - tz 1 * Spalte, letzter Absatz * Spalte, letzter Absatz * Spalte, Absatz 2 -	1-6,10	C12N5/06 G01N33/53 RECHERCHIERTE SACHGEBIETE (Int.Cl.7
Ole Reche in einem s der Techni Vollständig Unvollstän Nicht reche	LLSTÄNDIGE RECHEI rohenabtellung ist der Auffassung, de olohen Umfang nicht entspricht bzw. ik für diese Ansprüche nicht, bzw. nur g recherchierte Patentansprüche: dig recherchierte Patentansprüche: erchierte Patentansprüche: serchierte Patentansprüche: EBE Beschränkung der Recherche: E Ergänzungsblatt C	B ein oder mehrere Ansprüche, den Vorschrift entsprechen, daß sinnvolle Ermittlungen über d teilweise, möglich sind.	en dea EPÜ len Stand	GOIN
	Recharchenort MÜNCHEN TEGORIE DER GENANNTEN DOKU		unde liegende Ti	Prator buhr-Ebel, K heorien oder Grundsätze
X : von b Y : von b ander A : techn	esonderer Bedeutung allein betracht esonderer Bedeutung in Verbindung en Veröffentlichung derselben Kateg- ologischer Hintegrund schriftliche Offenbarung	E : alteres Patentidoki et nach dem Anmeldi mit einer D : in der Anmeldung one L : aus anderen Grün	ument, das jedoo edatum veröffent angeführtes Dok den angeführtes	h erst am oder licht worden ist ument Dokument



EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung EP 03 02 5506

	EINSCHLÄGIGE DOKUMENTE	,	KLASSIFIKATION DER ANMELDUNG (Int.CI.7)
Kategorie	Kennzeichnung des Dokuments mit Angabe, soweit erforderlich der maßgeblichen Teile	Betrifft Anspruch	
X	UCHIDA N ET AL: "Direct isolation of human central nervous system stem cells" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, Bd. 97, Nr. 26, 19. Dezember 2000 (2000-12-19), Seiten 14720-14725, XP002223508 ISSN: 0027-8424 * Zusammenfassung * * Seite 14722, rechte Spalte, letzter Absatz - Seite 14724, rechte Spalte, Absatz 1 * * Abbildungen 1,2 *	11-16,19	RECHERCHIERTE
X	KANEKO Y ET AL: "MUSASHI1: AN EVOLUTIONALLY CONSERVED MARKER FOR CNS PROGENITOR CELLS INCLUDING NEURAL STEM CELLS" DEVELOPMENTAL NEUROSCIENCE, S. KARGER, BASEL, CH, Bd. 22, Nr. 1/2, 2000, Seiten 139-153, XP001033925 ISSN: 0378-5866	11-16,19	SACHGEBIETE (Int.Cl.7
х	* Zusammenfassung * * Abbildung 5 * EP 1 354 943 A (NAT INST OF ADVANCED IND SCIEN) 22. Oktober 2003 (2003-10-22) "Monoclonal antibodies, hybridomas, cell isolation method, isolated cells and immunological diagnostic method" * Spalte 2, Zeile 42 - Spalte 3, Zeile 11 * * Spalte 13, Zeile 41 - Spalte 14, Zeile 53 *	1-19	



EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung EP 03 02 5506

	EINSCHLÄGIGE DOKUMENTE	KLASSIFIKATION DER ANMELDUNG (Int.Cl.7)	
Categorie	Kennzeichnung des Dokuments mit Angabe, soweit erforderlich der maßgeblichen Teile	Betrifft Anspruch	
×	GIMONA MARIO ET AL: "Beta-Actin Specific Monoclonal Antibody" CELL MOTILITY AND THE CYTOSKELETON, Bd. 27, Nr. 2, 1994, Seiten 108-116, XP009028901 ISSN: 0886-1544 * das ganze Dokument *	8,9,17,	
			RECHERCHIERTE SACHGEBIETE (Int.Cl.7)
	·		·

EPO FORM 1503 03.82 (POLC12)



UNVOLLSTÄNDIGE RECHERCHE ERGÄNZUNGSBLATT C

Nummer der Anmeidung EP 03 02 5506

Unvollständig recherchierte Ansprüche: 6, 16

Grund für die Beschränkung der Recherche (nicht patentfähige Erfindung(en)):

Artikel 52 (4) EPÜ – Verfahren zur chirurgischen Behandlung des menschlichen oder tierischen Körpers

Weitere Beschränkung der Recherche

Unvollständig recherchierte Ansprüche: 1-5, 7-15, 17-19

Grund für die Beschränkung der Recherche:

In den Listen A-E, auf die sich in den Patentansprüchen bezogen wird, sind insgesamt etwa 1000 putative Positiv- und Negativmarker neuraler Vorläuferzellen und neuraler Stammzellen aufgelistet. Diese putativen Marker sind teilweise bereits bekannte Proteine, wie z.B. beta-Aktin oder Interleukin 1 alpha, teilweise aber auch undefinierte, als "ESTs" benannte sogenannte Marker oder partielle mRNA-Sequenzen. Aufgrund der grossen Anzahl der putativen Marker und deren tw. mangelhaften Identifikation ist es unmöglich, eine vollständige Recherche zu erstellen.

ANHANG ZUM EUROPÄISCHEN RECHERCHENBERICHT ÜBER DIE EUROPÄISCHE PATENTANMELDUNG NR.

EP 03 02 5506

In diesem Anhang sind die Mitglieder der Patentfamilien der im obengenannten europäischen Recherchenbericht angeführten Patentdokumente angegeben.
Die Angaben über die Familien mitglieder entsprechen dem Stand der Datei des Europäischen Patentamts am Diese Angaben dienen nur zur Unterrichtung und erfolgen ohne Gewähr.

07-04-2004

Im Recherchenbericht angeführtes Patentdokument		Datum der Mitglied(er Veröffentlichung Patentfan		der lie	Datum der Veröffentlichung	
EP 1354943	A	22-10-2003	EP JP US	1354943 2004002350 2003186335	Α	22-10-2003 08-01-2004 02-10-2003

Für nähere Einzelheiten zu diesem Anhang : siehe Amtsblatt des Europäischen Patentamts, Nr.12/82